

EVALUATION OF HOUSEHOLDS' PRACTICES TOWARDS SOLID WASTE MANAGEMENT

MOHAMED OSMAN ELAMIN^{*1}, MARY ACHOL WILLIAM², HATIM RAHAMTALLA²,
ALI ALSHEHRI³ AND FOWZI ELAMIN⁴

^{1,3,4}*Faculty of Public Health & Informatics, Umm-Al-Qura University, KSA, Saudi Arabia*

²*Al-Zaiem AL-Azahari University, ³University of Bahri, Sudan*

(Received 24 October, 2021; Accepted 15 December, 2021)

ABSTRACT

Solid waste is the organic and inorganic waste materials such as product packaging, furniture, clothing, bottles, kitchen refuse, paper, paint cans, batteries, etc. Management of solid waste; is the process of collection, treatment, and recycling of solid waste in a sustainable manner to avoid the adverse effect on the environment. In Sudan waste management is poor and solid wastes are dumped along roadsides and into open areas, endangering health and attracting vermin. This descriptive cross-sectional community-based study with the aim to assess the practices of the households towards the management of house refuse was conducted in Al-Ozozab, Khartoum, 2019. Four hundred households were selected by multi stage sampling techniques from the community and data were collected by questionnaire and observation checklist. Most types (83.3%) of containers used for the waste store were plastic bags. All (100%) of the respondents cover the waste container. Almost all the respondents wash their hands with water and soaps after cleaning. 50.7% of respondents hear about hazardous domestic waste from TV and (19.3%) from radio. More than half (53.3%) of the houses' level of cleanliness was very good. Factors such as age, marital status, education level, family size, and monthly income significantly influence waste management practice. It can be concluded that the household has good level of practice towards solid waste management.

KEY WORDS: Evaluation, Households, Practices, Management, Solid Waste

INTRODUCTION

Solid waste is made up of organic and inorganic waste materials that comes about as a result of human and animal activities and is no longer needed which needs to be discarded due to its value loss to the user. It includes; product packaging, grass clippings, furniture, clothing, bottles, kitchen refuse, paper, appliances, paint cans, batteries, etc. These wastes are produced in the society and generally do not carry any value to its first user, (Vickey, 2019). The term solid waste management is the process of collection, treatment and recycling of solid waste in a sustainable manner to avoid the adverse effect on the environment, (Wilson, 2015). The first step of waste management is waste collection. House-to-house collection is very common in most developed

nations but very low in developing nations due to several challenges including financial, population expansion and other economic difficulties, (Bezama and Agamuthu, 2019). There are various sources of solid waste; they include household waste, industrial, commercial, construction and demolition, treatment plants and sites, agriculture and medical waste, (Paes, 2019). The houses are one of the major sources, which discard domestic solid waste materials daily, these garbage and wastes are; food residues, glass, vegetables, fruits, wood, papers, minerals and plastic. Therefore, segregation and transporting of these materials to the final disposal sites are important for the protecting human health and environment. Illegal waste dumping of solid waste raises many problems even in developed countries, transition economies, or developing

countries, where rural areas are frequently exposed to such environmental threats, (Zeng, 2015). Residents in developing countries, especially the urban poor, are more severely impacted by unsustainably managed waste. In low-income countries, over 90% of waste is often disposed in unregulated dumps or openly burned. These practices create serious health, safety, and environmental consequences. Poorly managed waste serves as a breeding ground for disease vectors, contributes to global climate change through methane generation, and can even promote urban violence (The World Bank, 2019).

For effective waste management, waste minimization, reuse, recycle and energy recovery are more sustainable than conventional landfill or dumpsite disposal technique. These actions not only help in protecting environment, but also help in employment generation and boosting up the economy, (Salman, 2019). Solid waste management techniques increase the rate of recycling waste and minimized waste disposal on land. Creative recycling is good for the environment, saves money, and can form the basis of an enjoyable and engaging hobby, (Hofverberg and Maivorsdotter, 2018).

Low and middle income countries face major challenges in ensuring universal access to waste collection services, eliminating uncontrolled disposal and burning and moving towards environmentally sound management for all wastes, (UNEP, 2015). Globally, 39% percent of the world population does not use proper method for waste disposal. Some 1.1 billion people still disposed waste in the open places. Improper waste disposal is most widely practiced in rural areas so that people face many health related problems, (Shahzadi, 2018).

Communities in developing countries often turn to waste disposal methods that have proven to be destructive to human health and the environment, such as open dumping and burning because they feel they have no other options to manage their solid waste, (Al-Khatib, 2015). This study aimed to assess the practices of the households towards the management of house refuse, Al-Ozozab, Khartoum, Sudan, 2019 and specifically to identify the practices concerning collection of house refuse, to determine the practices concerning transporting of house refuse and to explore the contributing factors of management of house refuse. The average weight of solid waste generated in Alkalakla Administrative Unit was 0.401 kg/ capita/day, and accordingly estimated annual amount of solid waste

was (36241.6 ton) (Elzaki and Elhassan, 2018).

Kiran stated that knowledge of often household removed their solid wastes and general solid wastes from the house were regularly (82.5%). He found that the majority of participants had a positive attitude towards solid waste disposal and 98.3% felt that improper solid waste removal and disposal effects environment. Regarding household waste disposal practice it was found to be unsatisfactory as 78 households disposed household's wastes by just throwing away outside the house, (Kiran *et al*, 2015). Barloa conducted a study to establish the effect of knowledge, attitudes and practices on waste management among 2528 Polytechnic university students. The findings indicate that 73.4% of the students' knowledge to be satisfactory, 71.4% had positive attitude towards strategic waste management issues; while around 43.1% depicted satisfactory levels in practice, (Barloa, 2016). Study on attitude of household towards waste management in a rural area of Northern Kerala revealed that the participants responsible for the waste management in household were women. The majority (82%) had an educational qualification of high school and above. Most of them (82.5%) were housewives. Almost 70% had the belief that government is not doing anything to fix the garbage problem. About 97%, 88.6% and 92% were willing to do composting, segregation and recycling of waste respectively. Majority of the participants had above average attitude towards household waste management, (Kaithery and Karunakaran, 2019). A study conducted by Putnick and Bornstein showed that girls are more likely to be involved in excessive housework than boys (Putnick and Bornstein, 2016). A study conducted by Laurieri, regarding the practical management of home waste, showed that respondents differentially deliver their small bins outside their house during the week according to the pick-up planning scheduled by municipal authorities, with only a very few people delivering their bins only 3 days a week or less (9.6%), (Laurieri, 2020). McAllister stated that the municipalities' responsibility is to organize and manage the public sanitation system, including providing the infrastructure for the collection, transportation, treatment and disposal of wastes (McAllister, 2015). A study on assessment of household waste management and hygienic practice was conducted in Ethiopia. The study discovered that (36%) of the households disposed solid wastes through municipality and 95.7% of households had

temporary storage means for solid waste. About 94.3% of the respondents revealed that the responsibility of waste management is left for women and girls. This study revealed that household management of waste in the community of Yirgalem town is poor in terms of their liquid waste management, (Tekleyohannes, 2019). A study conducted in Sierra Leone, showed that the solid waste generation and composition in Freetown was significantly affected by average family size, employment and marital status, monthly income, and number of room(s) occupied by households. He highlighted the role of socioeconomic factors in affecting the generation and composition of household solid waste, (Sankoh, 2015). A study that conducted in Kenya found that (65.3%) of the respondents mainly acquired the information on solid waste management through television while those in rural Gachororo acquired through radio (71.3%). Neighborhoods that are inhabited by affluent people may be more "cleaner" than those inhabited by low income groups due to the fact that those with higher monthly income are significantly more willing to contribute towards waste collection services than those with lower incomes, (King'oo, 2019). A study conducted in Shendi showed that 30% - 55% of the household produce about a kilogram a day per person, while 22% - 40% generate 5 or more kilograms daily. More than 50% of the household of different classes store their waste at home near the toilet, kitchen or other commodities in the house. Although most of the households were using suitable methods to store their solid waste (covered bins: 03%-09%; and closed plastic bags: 67%-74%), some household stored the waste at the collection point (03%-06%) or other improper methods (05%-18%), (Omer and Eltigani, 2018). Lee stated that participants washed their hands more frequently "after cleaning" (79.5%) (Lee, 2015).


METHODS AND MATERIALS

Study design: Cross sectional descriptive community based study with the aim to assess the practices of the households towards the management of solid waste, Al-Ozozab, Khartoum, Sudan, 2019. The study was conducted in Al-Ozozab area, Al-Shajara Locality, Khartoum State. The targeted population is households. The total number of households in the study area is 4577. 400 households were selected by the following formula:

$n = N/1+N (e^2)$. The sample was distributed by using systematic random sampling techniques. The data were collected by using questionnaire, interview and observation checklist. The collected data were analyzed by using Statistical Packages for Social Sciences (SPSS) version 20.0 and c2test was used to show the association between independent and dependent variables.

RESULTS


Nearly half (49.3%) of the of the respondents' age were less than 20 years, 48% were between 21 and 30 years old. The majority 98.5% of the respondents were married. 62% of the respondents were over secondary educated. Most of the respondents 71.8% were housewives. 53.5% of the respondents' monthly income ≥ 6000 SDGs, 21% between 4000



(n=400)

Fig. 1 Source of information about hazardous domestic waste


More than half (50.7%) of the respondents' source of information about hazardous domestic waste was TV followed by Radio (19.3%) and person (15.1%).



(n=400)

Fig. 2 Knowledge concerning how waste affects family members.

Nearly half (48.9%) of the respondents have good knowledge about waste affects health of the family members.



(n=400)

Fig. 3 Knowledge concerning the effect of domestic wastes on the environment

The majority (88.3%) of the respondents know the effect of domestic waste on the environment.

and 5000 SDGs, 16.3% between 3000-4000 SDGs and only 1% has less than 3000 SDGs.

Table 1 Attitudes towards hazard of domestic waste

Response	No.	%
Favorable	56	14.0
Unfavorable	344	86.0
Don't know	00	00
Total	400	100.0

(n=400)

The majority (86.0%) of the respondents have unfavorable attitudes towards hazard of domestic waste

Table 2 Attitudes towards the correct method for waste disposal

Response	No.	%
Favorable attitudes	385	96.3
Unfavorable attitudes	15	3.8
Total	400	100.0

(n=400)

The majority 96.3 has favorable attitude towards the correct method for waste disposal

Table 3 Distribution of the respondents according to the practice of cleaning their houses

Response	No.	%
Cleaning	400	100.0
Not cleaning	0	0.0
Total	400	100.0

(n=400)

All the respondents were cleaning their houses.

Table 4 Distribution of the respondents according to the responsibility of cleaning homes

Response	No.	%
Mother	363	90.8
Both mother and father	37	9.2
Total	400	100.0

(n=400)


90.8% of the mothers were the first responsible of cleaning home followed by both mothers and fathers 9.3%.

Table 5 Distribution of the respondents according to their role in cleaning homes

Response	No.	%
Having role	393	98.3
Not having role	7	1.8
Total	400	100.0

(n=400)

The majority 98.3% of respondents have role in home cleaning



(n=400)

Fig. 4 Types of waste containers.

The majority (83.3%) of the respondents use bags for storing waste

Table 6 The practice of covering the waste

Response	No.	%
Covering	400	100.0
Not Covering	0	0.0
Total	400	100.0

(n=400)

All respondents (100%) cover the waste containers

Table 7 The practice of washing hands after cleaning

Response	No.	%
Washing with water and soaps	397	99.3
Washing with, water only	3	.7
Total	400	100.0

(n=400)


The vast majority (99.3%) of respondents are washing their hands with water and soaps after cleaning.

Table 8 The responsibility of transporting the waste

Response	No.	%
Governmental body	381	95.3
Private sector	7	1.8
Others	12	2.9
Total	388	100.0

(n=400)

95.3% of the waste was transporting by governmental bodies



(n=400)

Fig. 5 Distribution of the respondents according to the disposing of waste in case of the absent of the waste disposal vehicle

Nearly half 48% of the respondents dispose the waste in the street 32% dispose it in front of the house.

Table 9 Association between respondents' age and waste management practice

Cleanliness status	Age			Total
	< 20	21-30	> 30	
Clean	16 (100%)	185 (96.4%)	192 (100%)	393 (98.3%)
Not clean	0 (0.0%)	7 (3.6%)	0 (0.0%)	7 (1.8%)

(n=400)

There is a significant associations between respondents' age and waste management practiceat (p<.05)

Table 10 Association between marital status and practice of hand washing

Hand washing	Marital status		Total
	Married	Not married	
Washing hands with water and soaps	392 (99.5%)	5 (83.3%)	397 (99.3%)
Washing hands with water only	2 (0.5%)	1 (16.7%)	3 (0.8%)

(n=400)

There was a significant associations between marital status and hand washing after cleaningat (p<.05)

Table 11 Association between the respondents' occupation and the frequency of cleaning the house

	Occupation			Total
	Housewife	Worker	Employee	
Twice daily	34 (11.8%)	24 (61.5%)	22 (29.7%)	80 (20%)
Once a day	229 (79.8%)	15 (38.5%)	50 (67.6%)	294 (73.5%)
Every two days	24 (8.4%)	0 (0.0%)	2 (2.7%)	26 (6.5%)

(n=400)

There was a significant association between the respondents' occupation and frequency of cleaning the houseat (p<.05)

DISCUSSION

The study illustrated that the majority of the mothers (90.8%) were the first responsible of cleaning home followed by both mother and father. This was due to the gender role of the culture and social norms of the community. This finding agreed with the study that was conducted in Ethiopia, which showed that 94.3% of the mothers and girls were the first responsible of cleaning home, (Tekleyohannes, 2019). The study indicated that the majority (98.3%) of respondents mentioned that boys and girls have a role in home cleaning. This indicated participation of both genders in home cleaning. This disagreed with the study that was conducted by Putnick and Bornstein, which showed that girls are more likely to be involved in excessive housework than boys, (Putnick and Bornstein, 2016). The current study showed that most types (83.3%) of containers used for storing waste were bags and these bags were covered. This because the plastic bags were available and less expensive compared with other types of waste containers. This finding agreed with a study that was conducted in Shendi

City, which found that most of the households were using plastic bags and 67%-74%, of the respondents covered the waste containers, (Omar and Eltigani, 2018). The study indicated that the vast majority (99.3%) of respondents wash their hands with water and soaps after cleaning. This due to the convinced messages from T.V as indicated in the study that 50.7% of the respondents received their health messages from T.V. This finding agreed with Lee study, who stated that participants washed their hands more frequently "after cleaning" (79.5%), (Lee *et al.*, 2015). Although, (86.0%) and (96.3%) of respondents have favorable attitudes regarding hazardous domestic waste and correct method for waste disposal, the majority (93%) place all the different types of waste in one container. This indicated that no segregation was done regarding solid waste at household level. This agreed with the study that was conducted in a rural area of Northern Kerala, which showed 93.8% of the study population had above average attitude and 6.2% had below average attitude but disagreed with the same study that found about 97%, 88.6% and 92% were willing to do composting, segregation and

recycling of waste respectively, (Kaithery and Karunakaran, 2019). The majority (98.2%) of the respondents mentioned that the government transports the waste. This finding indicates that the responsibility of waste transport depends on the government. This agreed with the study, which showed that the municipalities (government) have been in charge of providing solid waste management services in developing countries. However, the municipal responsibility is to organize and manage the public sanitation system, including providing the infrastructure for the collection, transportation, treatment and disposal of wastes (McAllister, 2015). 28.8% of the respondents reported that the frequent transporting of the waste was one per week. This may return to the lack of vehicles or trucks that transport the waste in the study area. This finding agreed with the study that was conducted by Laurieri, who found that the respondents differentially deliver their small bins outside their house during the week according to the pick-up planning scheduled by municipal authorities, with only a very few people delivering their bins only 3 days a week or less (9.6%), (Laurieri *et al.*, 2020). In case the waste vehicle does not attend, half of the respondents (48%) throw their solid waste in the street, 32% throw in front of the home. This may be due to poor practice concerning waste disposal. This agreed with the study that conducted by Kiran, who stated that household waste disposal practice was found to be unsatisfactory as (65%) households disposed of household wastes by just throw away outside the house (Kiran *et al.*, 2015). The current study showed significant association between respondents' age and cleaning solid waste. This implies that younger households were more likely to practice proper solid waste management compared to older ones. This agreed with study that conducted by Barloa, which shows that age and education levels were important contributing factors, (Barloa *et al.*, 2016). The study showed significant association between occupation and frequency of clean home. Housewives significantly clean their home once a day, followed by employee, twice a day. This may be due to spare time of the housewives compared with the employees. The finding agreed with the study that conducted by Sanko, who found an association between employment status and waste management, (Sankoh *et al.*, 2015). The study showed that those who have family income less than 3000 SDG were significantly often clean their kitchen twice (100%).

The finding indicated that income was not a major contributing factor in influencing solid waste management. This disagreed with Kingoo's statement that neighborhoods that are inhabited by affluent people may be more "cleaner" than those inhabited by low income groups due to the fact that those with higher monthly income are significantly more willing to contribute towards waste collection services than those with lower incomes, (Kingoo, 2019).

CONCLUSION

The study concluded that the majority 77.6% of the solid wastes generated at home were papers and trees papers, 7.5% rubble, 3% glass or iron, and 11.9% were animal waste. 87.3% of the solid waste was stored covered in tied plastic bags containers and 48% of the respondents dispose the waste in the street 32% dispose it in front of the house without sorting. Although, (97%) of the waste was transported with governmental body and disposed appropriately at communal sites, (32%) of the households members throw waste in any available space including front of houses and (48%) throw it in streets. (90.8%) of the mothers are responsible for household waste management. The TV represents (50.7%) of the solid waste management source of information and Radio represents (19.3%). Factors such as age, marital status, education level, family size and monthly income significantly influence waste management practices.

Recommendations

The State Ministry of Health should involve community members in decision-making and make use of the local available mass media (radios, televisions, newspapers, posters, magazines) to promote the practice of the households regarding solid waste management beside providing more communal trash bins and the Ministry of Education should include the management of solid waste in the outclass activities.

REFERENCES

- Al-Khatib, I. A., Kontogianni, S., Abu Nabaa, H., Alshami, N. and Al-Sari', M. I. 2015. Public perception of hazardousness caused by current trends of municipal solid waste management. *Waste Management*. 36323-36330.
- Bezama, A. and Agamuthu, P. 2019. *Addressing the Big Issues in Waste Management*. SAGE Publications,

- UK, London, England.
- Barloa, E. P., Lapie, L. P. and de la Cruz, C. P. P. 2016. Knowledge, attitudes, and practices on solid waste management among undergraduate students in a Philippine State University. *Journal of Environment and Earth Science*. 6(6): 146-153.
- Elzaki, A. A. M. and Elhassan, B. M. 2018. International Journal of Waste Resources.
- Hofverberg, H. and Maivorsdotter, N. 2018. Recycling, crafting and learning-An empirical analysis of how students learn with garments and textile refuse in a school remake project. Environmental Education Research.
- Kiran, K., Kini, S., Santhosh, N. and Kiran, N.U. 2015. KAP study of solid waste disposal of households in Kuttar and Manjanadi Panchayath. *Nitte University Journal of Health Science*. 5(3).
- Kaithery, N. N. and Karunakaran, U. 2019. Study on attitude of household waste management in a rural area of Northern Kerala. *International Journal of Community Medicine and Public Health*. 6(5): 2095.
- King'oo, S. N. 2019. *Assessment of household domestic waste management practices, Kiambu county-Kenya* (Doctoral dissertation).
- Lee, M. S., Hong, S. J. and Kim, Y. T. 2015. Hand washing with soap and national hand washing projects in Korea: 2006-2014. *Epidemiology and health*. 37.
- Laurieri, N., Lucchese, A., Marino, A. and Digiesi, S. 2020. A Door-to-Door Waste Collection System Case Study: A Survey on its Sustainability and Effectiveness. 12(14): 5520.
- McAllister, J. 2015. Factors influencing solid-waste management in the developing world.
- Omar Yousof, M. A. and Eltigani Osman, M. O. 2018. Assessment of municipal solid waste profile of Shendi City, River Nile State, Sudan.
- Putnick, D. L. and Bornstein, M. H. 2016. Girls and boys labor and household chores in low and middle income countries. *Monographs of the Society for Research in Child Development*. 81(1): 104.
- Paes, L. A. B., Bezerra, B. S., Deus, R. M., Jugend, D. and Battistelle, R. A. G. 2019. Organic solid waste management in a circular economy perspective. *Journal of Cleaner Production*.
- Shahzadi, A., Hussain, M., Afzal, M. and Gilani, S. A. 2018. Determination the level of knowledge, attitude, and practices regarding household waste disposal among people in rural community of Lahore. *International Journal of Social Sciences and Management*. 5(3).
- Sankoh, F. P., Yan, X. and Conteh, A. M. H. 2015. A situational assessment of socioeconomic factors affecting solid waste generation and composition in Freetown, Sierra Leone. *Journal of Environmental Protection*.
- Salman, Zafar, 2019. *Guide to Effective Waste Management*. Available at: <https://>
- Tekleyohannes, B. 2019. Assessment of Household Waste Management and Hygienic Practice in Yirgalem Town, Dale, Ethiopia. *Journal of Health and Environmental Research*. 5(2): 41-49.
- The World Bank 2019. Solid Waste Management. Available at: <https://www.worldbank.org/en/topic/urbandevelopment/brief/solid-waste-management>.
- UNEP, 2015. Global Waste Management Outlook, United Nations Environment Programme, Nairobi, Kenya.
- Vickey, 2019. What is Solid Waste - Sources and Classification? Available at: <https://civilengineeringnotes.com/solid-waste-classification>.
- Wilson, 2015. Global Waste Management Outlook. United Nations Environment programme Available online at: <http://www.eawag.ch/fileadmin/Do-main1/Abteilungen/sandec>.
- Zeng, C., Niu, D. and Zhao, Y. 2015. A comprehensive overview of rural solid waste management in China. *Frontiers of Environmental Science & Engineering*. 9(6): 949-961.