

## THE EFFECT OF AJWA DATES FRUIT EXTRACT ON OVARIAN FOLLICLE DEVELOPMENT IN FEMALE RATS EXPOSED TO ARSENIC

HERDHIKA A.R. KUSUMASARI<sup>1\*</sup>, HARWIN H DESYANTI<sup>1,2</sup>, MAHANUTABAH H. QURNIATILLAH<sup>1,3</sup> AND EVIANA NORAHMAWATI<sup>4</sup>

<sup>1</sup> Master of Midwifery Program, Brawijaya University, Malang, Indonesia

<sup>2</sup> Midwifery Program, Faculty of Health, Nurul Jadid University, Probolinggo, Indonesia

<sup>3</sup> Harapan Kita Women and Children Hospital, Jakarta, Indonesia

<sup>4</sup> Department of Anatomic Pathology, Faculty of Medicine, Brawijaya University, Malang, Indonesia

(Received 5 September, 2020; accepted 23 October, 2020)

### ABSTRACT

Arsenic trioxide (As<sub>2</sub>O<sub>3</sub>) is a form of inorganic arsenic which is very toxic in the environment. Arsenic exposure can cause reproductive health problems, one of which is the ovaries, by interfering with follicle development. Arsenic exposure causes oxidative stress in the ovaries due to the accumulation of excessive *Reactive Oxygen Species* (ROS) and an imbalance between ROS and antioxidants in the body. The powerful antioxidants contained in dates can against the effects of free radicals due to arsenic exposure. The group in this study was divided into a negative control group (without arsenic exposure), positive control (3 mg / kg BW arsenic), T1 (arsenic 3 mg / kg BW + date extract 2 ml/ kg BW), T2 (arsenic 3 mg / kg BW + date extract 4 ml/kg BW), and T3 (arsenic 3 mg/kg BW + date extract 8 ml/kg body weight). The administration of Ajwa date extract could significantly provide antioxidant protection in the treatment group with an increase in follicle count in this study.

**KEYWORD:** Arsenic trioxide, Arsen, Ovary, Free radical, *Reactive Oxygen Species*, Oxidative stress

### INTRODUCTION

Exposure to heavy metals, one of which is arsenic, which is a group *metalloid*, can cause reproductive dysfunction in women (Mendola *et al.*, 2008), and residual heavy metal exposure in the body can affect women's reproductive health (Bloom *et al.*, 2011). Arsenic toxicity has become a significant concern due to increased contamination to this compound in the air, water, and soil (Flora, 2014). *The Agency for Toxic Substances and Disease Registry* United States (ATSDR) has designated arsenic as No. 1 on the Priority List of Hazardous Substances over the lead, mercury, and *Vinyl Chloride* (ATSDR, 2019) (Shen *et al.*, 2013). Currently, the number of humans exposed to arsenic through the air, food, water, and soil is increasing all the time. Arsenic toxic effects are very

common in developing countries. Studies on the toxicity of arsenic exposure in women have been carried out in several countries. In a study conducted on women in Taiwan by Lei *et al.* (2015), arsenic levels in infertile women were higher than in pregnant women. A case study conducted on women in China who were exposed to arsenic found that arsenic can cause ovarian insufficiency / *Premature Ovarian Insufficiency* (Pan *et al.*, 2020). A drastic decrease in the number of follicles due to environmental factors such as heavy metal exposure, namely arsenic, can result in early menopause (Kumar *et al.*, 2012).

The effect of arsenic exposure on the ovarian organs can cause damage to ovarian morphology while folliculogenesis affects follicular regression. Arsenic exposure causes a decrease in the number of

follicles including primordial, primary, secondary and *de Graaf* follicles and increases the number of atretic follicles (Chattopadhyay and Ghosh, 2010; Akram *et al.*, 2010; Mehta and Hundal, 2016; Khatun *et al.*, 2018; Yu *et al.*, 2019).

Many studies have examined the benefits of dates fruit for females' reproductive health because they are believed affect on oocyte quality, sperm and ovum interactions, implantation and early embryonic development through antioxidant mechanisms (Abdi *et al.*, 2017; Saryono *et al.*, 2018). The antioxidant properties of Ajwa dates are expected to suppress free radicals through *scavenger* pathway, which can reduce disease proliferation due to the strong antioxidant content in Ajwa dates (Ahmed *et al.*, 2016; Al-Yahya *et al.*, 2016). Therefore, in this study, we are interested in conducting studies on the effects of Ajwadate extract (*Phoenix dactylifera L.*) on the female reproductive system, especially on follicle development in female Wistar rats (*Rattus norvegicus*) and the lack of research on the effects of Ajwa date palm extract in preventing exposure to arsenic metal toxicity.

## MATERIALS AND METHODS

### Chemical

Arsenic powder ( $As_2O_3$ ) produced by LobaChemie as much as 500 mg was dissolved in 1000 ml *Normal Saline* 0.9% then stirred with a *magnetic stirrer* on heating at 50 °C for 3-4 hours to make the arsenic powder completely dissolve.

### Preparation of the Extract

Types of dates used for the study were 400 grams of Ajwa dates. Selection of dates by selecting old dates, not rotten, separated from the seeds, then thinly sliced and the flesh of the dates is taken. The extract making refers to previous research, namely research by Vayalil (2002) and Al-Rasheed *et al.*, (2015) with slight modifications. The flesh of the Ajwa dates that have been separated from the seeds, mashed with a *mortar* and *pestle*, then added 1200 ml of water (ratio 1: 3, g/ml). The thick date palm extract was centrifuged at 4 °C for 15 minutes at a speed of 10,000 g, then the supernatant was taken.

### Animals and Ethics Statement

Research procedures have been accepted by the Health Research Ethics Commission, Faculty of Medicine, University of Brawijaya. In this study, there were five treatment groups where there were 5

repetitions for each group. Negative control was rats given *Normal Saline* and not given date palm extract and arsenic, positive control rats exposed to arsenic 3 mg/kg BW/day, treatment 1, 2 and 3 rats given date extracts with 3 different doses (2 ml / KgBW / day; 4 ml/KgBB/day; 8ml/KgBB/day), then exposed to 3 mg/KgBB / day arsenic for 30 days. The research was conducted at the Laboratory of Bioscience, Anatomical Pathology and Biochemistry Laboratory, Faculty of Medicine, University of Brawijaya.

### Vaginal Smears Collection

After 30 days of treatment, a swab was performed to determine the rat estrous cycle. Mice in the proestrus phase were terminated. The swab samples were stained with *methylene blue* and observed the morphology of the epithelial cells under a microscope (Sezer *et al.*, 2020).

### Histological Examination

**Number of Follicles.** Ovarian samples were stained with *Hematoxylin Eosin (H&E)*, criteria for follicular classification based on Myers *et al.* (2004). The results of the staining of the sample were then viewed under a microscope *Dot slide Olympus XC 10* and the number of follicles was calculated *using Dot Slide software*.

### Data Analysis

Data were analyzed using software *SPSS 25.0*

## RESULTS AND DISCUSSION

Note: The mean number of ovarian follicles based on the treatment group. On the mean  $\pm$  SD, if it contains different letters, there is a significant difference ( $p \leq 0.05$ ) and if it contains the same letter, there is no significant difference ( $p \geq 0.05$ ).

The number of primordial follicles showed a significant result with a p-value = 0.043. There was a decrease in the number of primordial follicles in the arsenic-exposed only group ( $2.4 \pm 2.607^a$ ) compared to the healthy or negative control groups ( $2.6 \pm 1.949^a$ ), but the decrease was not statistically significant. In the group of mice given Ajwa date extract at a dose of 4 ml/kg BW, there was a significant increase in the number of primordial follicles compared to the positive control group that was given arsenic alone.

The same thing happened to the number of primary follicles, all treatment groups showed

significant results statistically with a value of  $p = 0.029$ . Although in the positive control group ( $2.4 \pm 2.607^a$ ) there was a decrease in the number of secondary follicles, the decline was not significant compared to the negative control group ( $2.6 \pm 1.949^a$ ) statistically. Ajwa date extract caused a protective effect on the number of primary follicles in female rats exposed to arsenic with the highest dose of date extract, 8 ml/kg BW. In treatment group 3 ( $3.0 \pm 1.000^{ab}$ ) there was a significant increase in the number of primary follicles compared to the arsenic exposed group ( $1.6 \pm 0.547^a$ ).

In the number of secondary follicles, the group of mice exposed to arsenic ( $1.2 \pm 1.095^a$ ) decreased in number but not significantly compared to the group of rats that were either healthy or not exposed to arsenic ( $2.4 \pm 1.140^a$ ). In the treatment group given date palm extract 8 ml/kg BW or Treatment 3 ( $2.0 \pm 0.707^a$ ) there was an increase in the number of secondary follicles compared to the positive control group ( $1.2 \pm 1.095^a$ ), the mean value of the P3 group approached the negative control group ( $2.4 \pm 1.140^a$ ). However, there was an increase in the number of secondary follicles with a mean value that exceeded the group of healthy rats that occurred in the treatment group given date extract 4 ml/kg BW ( $4.2 \pm 3.768^a$ ).

Then the number of early antral follicles also decreased the number of early antral follicles in the positive control group ( $0.8 \pm 0.836^a$ ) compared to the negative control group/only given Normal saline ( $1.2 \pm 0.447^a$ ), but the decrease was not significant. There were interesting results in the group of mice given date palm extract at a dose of 4 ml/kg BW ( $1.2 \pm 0.446^a$ ) and 8 ml/kg BW ( $1.2 \pm 0.447^a$ ) where the value of the number of early antral follicles increased, and the mean value was the same as healthy mice group ( $1.2 \pm 0.447^a$ ), whereas, at the smallest dose of date extract, 2 ml/kg BW, there was an increase in the number of early antral follicles ( $2.0 \pm 1.000^a$ ) compared to the mice exposed to arsenic alone.

In addition to the number of follicles mentioned in this study, researchers also counted the number of antral or *Graafian follicles* that was published. The results obtained were that there was a significant decrease in the number of *Graafian follicles* in mice exposed to arsenic and date palm extracts can increase the number of *Graafian follicles* in female rats exposed to arsenic (Kusumasari *et al.*, 2020).

The decrease in the number of follicles in the group of rats exposed to arsenic in this study was the same as previous research, namely Mehta and Hundal (2016) that rats exposed to arsenic caused problems in folliculogenesis. The decrease in the number of follicles occurs due to the mechanism of arsenic toxicity in various ways so that it affects the disruption of folliculogenesis. One of them is arsenic exposure causes an increase in *Reactive Oxygen Species (ROS)* which results in oxidative stress on the ovaries (Flora, 2011; Sun *et al.*, 2016). The increase in ROS is usually followed by an increase in lipid peroxidase and MDA which are biomarkers of oxidative stress. Homeostatic changes in the body, such as increased levels of ROS due to arsenic exposure, also alter the activity of antioxidant enzymes, which act as a first-line defence against free radicals. Inhibition of antioxidant enzymes will cause excessive levels of superoxide anions (Banerjee *et al.*, 2014), excessive accumulation of superoxide anions can cause follicular regression (Khatun *et al.*, 2018). Arsenic exposure also inhibits the peroxidase enzyme in the follicular fluid so that it interferes with folliculogenesis due to high levels of free radicals (Chattopadhyay and Ghosh, 2010). Apart from going through the ROS pathway, arsenic toxicity can damage DNA and damage ovarian cells (Akram *et al.*, 2009). A drastic decrease in the number of follicles caused by environmental factors such as heavy metal exposure, namely arsenic, can result in premature menopause in women (Kumar *et al.*, 2012).

The effect of arsenic on the endocrine system is also one of the causes of disturbances in follicular

**Table 1.** Effect of Ajwa Dates Extract on Ovarian Follicles Count of Female Rats.

Follicle	Negative Control Mean±SD	Positive Control (arsenic) Mean±SD	Treatment 1 (arsenic+date extract 2 ml/kgBB) Mean±SD	Treatment 2 (arsenic+date 4 ml/kgBB) Mean±SD	Treatment 3 (arsenic+date 8 ml/kgBB) Mean±SD
Primordial	2.6±1.949 <sup>a</sup>	2.4±2.607 <sup>a</sup>	3.0 ±2.645 <sup>a</sup>	7.0±3.082 <sup>ab</sup>	13.2±12.336 <sup>a</sup>
Primer	2.6±1.516 <sup>a</sup>	1.6±0.547 <sup>a</sup>	1.2 ±0.447 <sup>a</sup>	1.8±0.447 <sup>a</sup>	3.0±1.000 <sup>ab</sup>
Sekunder	2.4±1.140 <sup>a</sup>	1.2±1.095 <sup>a</sup>	1.2 ±0.836 <sup>a</sup>	4.2±3.768 <sup>a</sup>	2.0±0.707 <sup>a</sup>
Early Antral	1.2±0.447 <sup>a</sup>	0.8±0.836 <sup>a</sup>	2.0 ±1.000 <sup>a</sup>	1.2±0.446 <sup>a</sup>	1.2±0.447 <sup>a</sup>

development. Arsenic exposure indirectly interferes with the regulation of the ovarian pituitary axis by affecting the levels of the *Follicle Stimulating Hormone* (FSH) hormone and decreasing levels of the steroid estrogen hormone where the action of this hormone is needed in the folliculogenesis process. If there is inhibition of hormones that affect follicle development, it can result in low follicle regression and ovarian mass (Khatun *et al.*, 2018). Therefore, in this study, the group of mice exposed to arsenic alone had a decreased number of follicles than the healthy group of mice.

The preventive effect of Ajwa date fruit extract in this study is due to the antioxidant content contained in dates, namely melatonin, carotene, phenolic compounds, flavonoids and contains vitamins C, E and vitamin A. Phenolic compounds contained in dates are routine, catechin, *caffeic acid*, *gallic acid*, *quercetin*, *luteolin* and *apigenin* (Chaira *et al.*, 2007; Al-Farsi and Lee, 2008; Ben Abdallah *et al.*, 2009; Ali *et al.*, 2015; Eid *et al.*, 2015; Khalid *et al.*, 2017). Apart from phenolic compounds, a study conducted by Chao (2007) proved that dates contain estradiol.

Antioxidants in Ajwa date can inhibit lipid peroxidase and stop reactions associated with oxidative stress (Zhang *et al.*, 2013). The antioxidants contained in Ajwa dates can stop the bonding reactions associated with oxidative stress by reducing ROS formation (Ben Abdallah *et al.*, 2009). The phenolic content of Ajwa dates, with their redox properties, can act as a *scavenger* ROS, neutralize free radicals, and break down peroxides (Kchaou *et al.*, 2014). So that in this study proved that Ajwa date extract had a protective effect on ovarian follicles in female rats exposed to arsenic.

### CONCLUSION

This study shows that arsenic exposure can affect the number of ovarian follicles and Ajwa date extract, which contains high antioxidants that can protect ovarian follicles from damage caused by arsenic exposure.

### ACKNOWLEDGEMENT

Thanks to the laboratory analysts of Bioscience, Pathology Anatomy, Biochemistry and Laboratory of the Center for Biological Sciences (LSIH), Brawijaya University.

### REFERENCES

- Abdi, F., Roozbeh, N. and Mortazavian, A.M. 2017. Effects of date palm pollen on fertility: Research proposal for a systematic review. *BMC Research Notes*. 10(1) : 1-5.
- Ahmed, A. 2016. Nutritional probing and HPLC profiling of roasted date pit powder. *Pakistan Journal of Nutrition*. 15(3) : 229-237.
- Akram, Z. 2009. Genotoxicity of sodium arsenite and DNA fragmentation in ovarian cells of rat. *Toxicology Letters*. 190(1) : 81-85.
- Akram, Z. 2010. Adverse effects of arsenic exposure on uterine function and structure in female rat. *Experimental and Toxicologic Pathology*. 62(4): 451-459. Available at: <http://dx.doi.org/10.1016/j.etp.2009.07.008>.
- Al-Farsi, M.A. and Lee, C.Y. 2008. Optimization of phenolics and dietary fibre extraction from date seeds. *Food Chemistry*. 108(3) : 977-985.
- Ali, E., Sulieman, A.M.E., Elhafise, I.A.A. and Abdelrahim, A.M. 2015. Comparative Study on Five Sudanese Date (*Phoenix dactylifera* L.) Fruit Cultivars Comparative Study on Five Sudanese Date (*Phoenix dactylifera* L.) Fruit Cultivars., 2012(NOVEMBER) : 1245-1251.
- Al-rasheed, Nouf M. 2015. Aqueous Date Flesh or Pits Extract Attenuates Liver Fibrosis via Suppression of Hepatic Stellate Cell Activation and Reduction of Inflammatory Cytokines, Transforming Growth Factor- $\beta$ 1 and Angiogenic Markers in Carbon Tetrachloride-Intoxicated Rats. *Hindawi Publishing Corporation Evidence-Based Complementary and Alternative Medicine*, Volume 201, p.19 pages.
- Al-Yahya, M. 2016. 'Ajwa' dates (*Phoenix dactylifera* L.) extract ameliorates isoproterenol-induced cardiomyopathy through downregulation of oxidative, inflammatory and apoptotic molecules in rodent model. *Phytomedicine*. 23(11) : 1240-1248.
- Banerjee, Sayani, 2014. Female Reproductive Aging Is Master-Planned at the Level of Ovary Franks, S., (ed.). *PLoS ONE*. 9(5) : e96210. Available at: <https://dx.plos.org/10.1371/journal.pone.0096210> [Accessed: 16 January 2020].
- Ben Abdallah, F. 2009. Effects of date seed oil on testicular antioxidant enzymes and epididymal sperm characteristics in male mice. *Andrologia*. 41(4) : 229-234.
- Bloom, M.S. 2011. Associations between blood metals and fecundity among women residing in New York State. *Reproductive Toxicology*. 31(2) : 158-163. Available at: <http://dx.doi.org/10.1016/j.reprotox.2010.09.013>.
- Chaira, N., Ferchichi, A., Mrabet, A. and Sghairoun, M., 2007. Chemical Composition of the Flesg and the Pit of Date Palme Friut 2007.Pdf. *Pakistan Journal of Biological Science*. 10(13) : 2202-2207.

- Chao, C.T. and Krueger, R.R., 2007. The date palm (*Phoenix dactylifera* L.): overview of biology, uses, and cultivation. *Hort Science*. 42(5) : 1077-1082.
- Chattopadhyay, S. and Ghosh, D., 2010. Role of dietary GSH in the amelioration of sodium arsenite-induced ovarian and uterine disorders. *Reproductive Toxicology*. 30(3) : 481-488. Available at: <http://dx.doi.org/10.1016/j.reprotox.2010.05.002>.
- Chuan-Rui Zhang, Saleh A. Aldosari, Polana S. P. V. Vidyasagar, Karun, M. Nair, and M.G.N. 2013. Antioxidant and Anti-inflammatory Assays Con firm Bioactive Compounds in Ajwa Date Fruit. *Journal of Agricultural and Food Chemistry*.
- Eid, N. 2015. Impact of palm date consumption on microbiota growth and large intestinal health: A randomised, controlled, cross-over, human intervention study. *British Journal of Nutrition*, 114(8) : 1226-1236.
- Flora, S.J.S., 2011. Arsenic-induced oxidative stress and its reversibility. *Free Radical Biology and Medicine*, 51(2) : 257-281. Available at: <http://dx.doi.org/10.1016/j.freeradbiomed.2011.04.008>.
- Flora, S.J.S., 2014. *Handbook of Arsenic Toxicology*. Academic Press.
- Kchaou, W., Abbès, F., Attia, H. and Besbes, S. 2014. In vitro antioxidant activities of three selected dates from Tunisia (*Phoenix dactylifera* L.). *Journal of Chemistry*, 2014.
- Khalid, S. 2017. A review on chemistry and pharmacology of Ajwa date fruit and pit. *Trends in Food Science and Technology*. 63 : 60-69. Available at: <http://dx.doi.org/10.1016/j.tifs.2017.02.009>.
- Khatun, S. 2018. Spirulina platensis ameliorates arsenic-mediated uterine damage and ovarian steroidogenic disorder. *Facets*. 3 (1) : 736-753.
- Kumar, M. 2012. Chromosomal abnormalities and oxidative stress in women with premature ovarian failure (POF). *Indian Journal of Medical Research*. 135 (1) : 92-97.
- Kusumasari, H.A.R., Desyanti, H.H. and Mahanutabah, H. 2020. The Effect of Ajwa Dates Fruit Extract on Follicle Stimulating Hormone (FSH), Graafian Follicle and Endometrial Thickness in Female Rats Exposed to Arsenic. *Journal of Global Pharma Technology*. 12(08) : 122-130.
- Lei, H.L. 2015. Relationship between risk factors for infertility in women and lead, cadmium, and arsenic blood levels: A cross-sectional study from Taiwan. *BMC Public Health*. 15(1) : 1-11. Available at: <http://dx.doi.org/10.1186/s12889-015-2564-x>.
- Mehta, M. and Hundal, S.S. 2016. Effect of sodium arsenite on reproductive organs of female Wistar rats. *Archives of Environmental and Occupational Health*. 71 (1) : 16-25.
- Mendola, P., Messer, L.C. and Rappazzo, K. 2008. Science linking environmental contaminant exposures with fertility and reproductive health impacts in the adult female. *Fertility and Sterility*, 89(2 SUPPL.) : 81-94.
- Myers, M. 2004. Methods for quantifying follicular numbers within the mouse ovary. *Reproduction*. 127(5) : 569-580.
- Pan, W. 2020. A case-control study of arsenic exposure with the risk of primary ovarian insufficiency in women. *Environmental Science and Pollution Research*. 27(20) : 25220-25229.
- Saryono, S., Dwi, M. and Rahmawati, E., 2018. Effects of Dates Fruit (*Phoenix dactylifera* L.) in the Female Reproductive Process. 03(October), pp.1630-1633. Available at: [https://www.academia.edu/33897003/EFFECTS\\_OF\\_DATES\\_FRUIT\\_PHOENIX\\_DACTYLIFERA\\_L.\\_IN\\_THE\\_FEMALE\\_REPRODUCTIVE\\_PROCESS](https://www.academia.edu/33897003/EFFECTS_OF_DATES_FRUIT_PHOENIX_DACTYLIFERA_L._IN_THE_FEMALE_REPRODUCTIVE_PROCESS).
- Sezer, Z. 2020. Effects of vitamin E on nicotine-induced lipid peroxidation in rat granulosa cells: Folliculogenesis. *Reproductive Biology*. 20(1) : pp.63-74.
- Shen, S. 2013. Arsenic binding to proteins. *Chemical Reviews*. 113(10) : 7769-7792.
- Sun, H.J. 2016. Mechanisms of arsenic disruption on gonadal, adrenal and thyroid endocrine systems in humans: A review. *Environmental International*. 95 : 61-68. Available at: <http://dx.doi.org/10.1016/j.envint.2016.07.020>.
- Vayalil, P.K. 2002. Antioxidant and antimutagenic properties of aqueous extract of date fruit (*Phoenix dactylifera* L. Arecaceae). *Journal of Agricultural and Food Chemistry*. 50(3) : 610-617.
- Yu, H. 2019. Sodium Arsenite Injection Induces Ovarian Oxidative Stress and Affects Steroidogenesis in Rats. *Biological Trace Element Research*. 189 (1) : 186-193.